

Compactified Jacobian of some Unibranch curves

Jyotsna Gokhale
 Mehta Research Institute
 10 Kasturba Gandhi Marg
 Allahabad 211002
 India.
 E-mail : jgokhale@mri.ernet.in

Introduction

Let C be an integral curve with one unibranch singularity of embedding dimension greater than two. It is known that $(Pic^d C)^=$, the moduli of torsion free sheaves of rank one and degree d , is irreducible if and only if C is locally planar ([D], [AIK], [R], [K]). It is enough to consider a curve with one singularity [SK].

In this work ¹ we give a basic decomposition of $(Pic^0 C)^=$ towards irreducible components for an integral curve with one unibranch singularity, and give the decomposition into irreducible components for some types of curves. $\diamond$

¹ This work was done during '84 - '86 at Brandeis University as part of my doctoral dissertation, and for various reasons remained unpublished so far along with rest of my dissertation. It was brought to my notice in '92, when this paper was sent to Mathematische Zeitschrift for publication, that G.Pfister and J.H.M.Steenbrink have recently ('91) published a paper similar in content. We have consequently shortened the paper by giving the reference of Pfister and Steenbrink whenever possible, omitting repetition of already published proofs.

It is a duty and a pleasure to thank my advisor David Eisenbud for all the insight and the constructive comments he provided me with, and for the atmosphere he generates around him.

Notation :

k	an algebraically closed field of characteristic zero
C	an integral curve over k with one unibranch singularity Q
$C^\sim$	normalisation of C
π	the normalisation map
$\mathcal{O}$	completion of local ring of C at Q
$\mathcal{O}^\sim$	normalisation of $\mathcal{O}$
$\mathcal{M}$	maximal ideal of $\mathcal{O}$
$\mathcal{C}$	the conductor $Hom_{\mathcal{O}}(\mathcal{O}^\sim, \mathcal{O})$ of C
$\mathcal{M}_Q$	maximal ideal sheaf of Q
$\mathcal{C}_Q$	conductor sheaf
δ	$rk_k(\mathcal{O}^\sim/\mathcal{O})$; we omit the subscript k normally
$\mathcal{F}$	torsion free sheaf of rank one
$\mathcal{L}$	a locally free sheaf of rank 1
$\mathcal{W}_C^0$	the canonical dualising sheaf of C
C'	first partial normalization of C such that $rk_k(\mathcal{O}'/\mathcal{O}) = 1$ and $C' \neq \mathcal{C}$; i.e., $\mathcal{O}' = \mathcal{O} + t^{-1}\mathcal{C}$.
C^i	successive partial normalizations defined along the same line
$\mathcal{M}^i$	maximal ideal of $\mathcal{O}^i$
$\mathcal{C}^i$	the conductor of C^i
$Pic^0 C$	Picard group of degree 0 of C
π^*	the map $Pic^0 C \longrightarrow Pic^0 C^\sim$ of Picard groups corresponding to π
K	$ker(\pi^*)$
$(Pic^0 C)^\sim$	moduli of torsion free sheaves of degree 0 and rank one.
$\overline{S}$	Zariski closure of a variety or scheme S
$\overline{(Pic^0 C)}$	Zariski closure of $Pic^0 C$ in $(Pic^0 C)^\sim$
$B(\overline{Pic^0 C})$	boundary of $\overline{Pic^0 C}$, def(0.1)
$\chi(\mathcal{F})$	Euler characteristic of $\mathcal{F}$
$deg(\mathcal{F})$	$\chi(\mathcal{F}) - \chi(\mathcal{O}_C)$
Γ	$\Gamma(C) = (k_1, \dots, k_r) \mathbf{N}$, the semigroup of orders of generators of $\mathcal{O}$ over k , assuming $2 < k_1 < k_2 < \dots < k_r$
v_0	order of the conductor, i.e., $\mathcal{C} = t^{v_0}\mathcal{O}^\sim$
t	any local parameter of $\mathcal{O}^\sim$. ◇

2

² Γ^* of [PS] corresponds to our C' .

0: Preliminaries

We define $(Pic_C^d)^=(S) = \{ \text{isomorphism classes of torsion free } \mathcal{O}_C(S) \text{-modules of degree } d \text{ and rank } 1 \}$. $(Pic_C^d)^=$ is representable and is represented by a projective scheme $(Pic^d C)^=$ [D], [AK], which is irreducible [R], [K], iff $rk(\mathcal{M}/\mathcal{M}^2) \leq 2$. We define the boundary of $\overline{Pic C}$, a la [R]:

Definition 0.1: $\mathcal{F} \in B(\overline{Pic^0 C})$ if $\mathcal{F} \notin Pic^0 C$, there exists an $\mathcal{O}_{C \times_{Spec k[[t]]} Spec k[[t]]}$ -module $\mathcal{G}$ flat over $k[[t]]$ such that $(\mathcal{G}/t\mathcal{G}) \approx \mathcal{F}$ and the stalk of $\mathcal{G}$ at the generic point of $Spec k[[t]]$ is a locally free $\mathcal{O}_C$ -module. $\parallel$

Let $K = ker(\pi^*)$, where $\pi^* : Pic^0 C \longrightarrow Pic^0 C^\sim$ is the map of respective Picard groups corresponding to the normalization map $\pi : C^\sim \longrightarrow C$.

[R,1.2]: $\mathcal{F} \in \overline{Pic^0 C} \iff \exists \mathcal{L} \in Pic^0 C$ such that $\mathcal{F} \otimes \mathcal{L} \in \overline{K}$. $\parallel$

For $d \leq rk(\mathcal{O}^\sim/C)$, the functor

$$E(\mathcal{C}, d)(S) = \{ \text{isomorphism classes of } \mathcal{O}_S\text{-modules } F_S \mid \mathcal{C} \otimes_k \mathcal{O}_S \subseteq F_S \subseteq \mathcal{O}^\sim \otimes \mathcal{O}_S, \text{ and } rk(\mathcal{O}^\sim \otimes \mathcal{O}_S/F_S) = d \}$$

is representable by a projective scheme [R], also called $E(\mathcal{C}, d)$; we identify $E(\mathcal{C}, d)$ with its image in $(Pic^0 C)^=$. In particular, since $K \subseteq E(\mathcal{C}, \delta)$, every boundary point “defines an element of $E(\mathcal{C}, \delta)$ ” [R, Th.2.3(b)]. So it is enough to consider limits of one - parameter families of free $\mathcal{O}$ -modules $\{F_\beta\} \subseteq E(\mathcal{C}, \delta)$ parameterized by $k[[\beta]]$. A free module $\mathcal{G} \in E(\mathcal{C}, \delta)(k(\beta))$ is necessarily of the kind $\mathcal{G} = \partial_\beta \mathcal{O}$, $\partial_\beta \in [\mathcal{O} \otimes k(\beta)]^*$. We shall write $\partial_\beta \mathcal{O} \longrightarrow F$ if F is in the closure of such a family.

A key concept of help in determining $B(\overline{Pic^0 C})$ from [J] is that of the unique filtration of $\mathcal{O}$ given by $\mathcal{O} \supseteq \mathcal{M} \supseteq \dots \supseteq I_j \dots \supseteq \mathcal{C}$ such that $rk_k(I_j/I_{j+1}) = 1$, and $ord_t(I_j) < ord_t(I_{j+1}) \ \forall j \geq 0$.³ Every $F \in E(\mathcal{C}, \delta)$ has such a unique filtration. We define [J'] a functor $Filt_{\mathcal{C}}^\delta$ using this concept of filtrations of $\mathcal{F}$ and proved that it is representable by a projective scheme $Filt(\mathcal{C}, \delta) = \{F \in E(\mathcal{C}, \delta) \mid F_j \subseteq I_j \mathcal{O}^\sim \ \forall j \geq 0\}$. Since $\overline{K} \subseteq Filt(\mathcal{C}, \delta) \subseteq E(\mathcal{C}, \delta)$ this is a major tool in determining $B(\overline{Pic^0 C})$. For brevity we shall use the notation $Filt(\mathcal{C}, \delta)$ in this work.

We will close the section with some remarks implicitly assumed and used in further sections. Since $v_0 - \delta$ is used frequently, we will use notation $v_0 - \delta = \gamma$ for ease of reading.

Remarks:

(0.2) Since $\#\{j \in \mathbf{N} \mid j \notin \Gamma\} = \delta$, so $\#\{j \in \Gamma \mid j < v_0\} = \gamma$.

³The definition of I_n in [PS] is parallel to this.

(0.3) $j \in \Gamma \implies v_0 - 1 - j \notin \Gamma \ \forall j \in \mathbf{Z}$. The converse is true iff C is Gorenstein [HK].

(0.4) (i) $\delta - 1 \leq v_0 \leq 2\delta$.
(ii) $\#\{j \in \mathbf{N} \mid j, v_0 - 1 - j \notin \Gamma\} = 2\delta - v_0$.

(0.5) If F is an $\mathcal{O}$ -module such that $\mathcal{C} \subseteq F \subseteq \mathcal{O}^\sim$, and $\text{End}(F)$ does not contain $t^{-1}\mathcal{O}$, then $\mathcal{O} \subset F \subseteq \mathcal{O}^\sim$, and $v_0 - \delta \leq rk(F/\mathcal{C}) \leq \delta$. Further, if $rk(F/\mathcal{C}) = \delta$ then $F \approx \mathcal{W}_{C,Q}^0$ necessarily. Conversely, for $j : v_0 - \delta \leq j \leq \delta$, there exists an F such that $\mathcal{C} \subseteq F \subseteq \mathcal{O}^\sim$, $rk(F/\mathcal{C}) = j$, and $\text{End}(F)$ does not contain $t^{-1}\mathcal{C}$.

(0.6) The following are equivalent:

- (i) $v_0 = k_{\gamma-1} + k_1$.
- (ii) $k_i = ik_1 \ \forall i \leq \gamma, \ k_i \in \Gamma$.
- (iii) $rk(\mathcal{M}/\mathcal{M}^2 + t\mathcal{C}) = 1$.

(0.7) $E(\mathcal{C}, \delta)$ is irreducible $\iff \overline{K} = E(\mathcal{C}, \delta)$.

(0.8) If $\text{Hom}_{\mathcal{O}_c}(\mathcal{F}, \mathcal{O}_c) = \mathcal{F}^v \in B(\overline{\text{Pic}^0 C})$ then $F \in B(\overline{\text{Pic}^0 C})$.

(0.9) $I_j \in \overline{K} \ \forall j \geq 0$, since $I_j = \text{Hom}_{\mathcal{O}}(\mathcal{O}^j, \mathcal{O}) = (\mathcal{O}^i)^v$. In particular $\mathcal{M}_Q \in \overline{\text{Pic}^0 C}$.

(0.10) $\pi'_*(\overline{\text{Pic}^{-1} C'}) \subseteq B(\overline{\text{Pic}^0 C})$ [J,(1.1)].

(0.11) $B(\overline{\text{Pic}^0 C}) \subseteq \pi'_*(\text{Pic}^{-1} C)^\#$ [J,(1.2)].

(0.12) $B(\overline{\text{Pic}^0 C}) = \bigcup_i \pi'_*(\overline{\text{Pic}^{-i} C^i}) \iff \mathcal{M} = \mathcal{C}$ [J,(1.3)].

(0.13) The following conditions are equivalent [J, §2] :

- (i) $rk(\mathcal{M}/\mathcal{M}^2 + t\mathcal{C}) = 1$
- (ii) $B(\overline{\text{Pic}^0 C}) = \pi'_*\overline{\text{Pic}^{-1} C'}$
- (iii) $E(\mathcal{C}, \delta) = \overline{K}$ and $E(\mathcal{C}', \delta - 1) = \overline{K'}$.

(0.14) If $B(\overline{\text{Pic}^0 C}) = \pi'_*(\text{Pic}^{-1} C')^\#$ then either C is locally planar or $\text{ord}_t \mathcal{C} = v_0 = 2\delta - 1$ [J,§2]. $\diamond$

1: Decomposition.

In this section we give the basic decomposition of $(\text{Pic}^0 C)^\#$ towards irreducible components. If F is an $\mathcal{O}$ -module such that $\mathcal{O} \subseteq F \subseteq \mathcal{O}^\sim$, and $\text{End}(F)$ does not contain $\mathcal{O}'$, then $F' = uF$, $u \in \mathcal{O}^{\sim*}$ for any $F' \approx F$ such that $\mathcal{C} \subseteq F' \subseteq \mathcal{O}^\sim$. In particular $rk(F'/\mathcal{C}) = rk(F/\mathcal{C})$ is fixed. For

r such that $v_0 - \delta - 1 \leq r \leq \delta - 1$, we define ⁴

Definition (1.0): $Pic_r^0 C = \{\mathcal{F} \in (Pic^0 C)^\sim \mid rk(\mathcal{F}/\mathcal{F}\mathcal{C}) = r + 1\}$.
Equivalently, $Pic_r^0 C = \{\mathcal{F} \in (Pic^0 C)^\sim \mid \exists F \approx \mathcal{F}_Q \text{ such that } \mathcal{O} \subseteq F \subseteq \mathcal{O}^\sim, \text{ and } rk(F/\mathcal{C}) = r + 1\}$.

Lemma 1.1: From definition of $Pic_r^0 C$,

- (i) $Pic_r^0 C \cap Pic_{r'}^0 C$ is empty for $r \neq r'$.
- (ii) $(Pic^0 C)^\sim = \bigcup \{Pic_r^0 C \mid r = \gamma - 1, \dots, v_0 - 2\}$. ⁵
- (iii) $\overline{Pic_r^0 C} \subseteq Orb_{Pic^0 C} E(\mathcal{C}, \delta - r) = \{\mathcal{F} \in (Pic^0 C)^\sim \mid \exists F \approx \mathcal{F}_Q \text{ such that } \mathcal{C} \subseteq F \subseteq \mathcal{O}^\sim \text{ and } rk(F/\mathcal{C}) = r + 1\}$. ||

We would like to know if and when $\overline{Pic_r^0 C}$ is a component of $(Pic^0 C)^\sim$. Generally $\overline{Pic_r^0 C}$ is not irreducible. However, since for all r in this range $\exists \mathcal{F} \in Pic_r^0 C$ such that $End(\mathcal{F})$ does not contain $\mathcal{O}_{C'}$, ⁶

Lemma 1.2: $\overline{Pic_r^0 C}$ contains an irreducible component of $(Pic^0 C)^\sim$ for r such that $\gamma - 2 < r < \delta$. ||

Therefore

Theorem 1.3: The minimum number of components of $Pic^0 C$ for a general unibranch singularity is $2\delta - v_0 + 1$, and these are the components containing an open subvariety $\{\mathcal{F} \mid End(\mathcal{F}) \text{ does not contain } \mathcal{O}_{C'}\}$. ||

This bound is strict for some types of curves such as one with a non Gorenstein singularity of embedding dimension greater than two, and becomes an equality for locally planar or $\mathcal{M} = \mathcal{C}$ type singularities.

We generalize [R,1.2]: let $(-)^*$ denote $Hom(-, \mathcal{O}_{C^\sim})$ and let $K_r = ker(\pi_r^*)$ where $\pi_r^*: Pic_r^0 \longrightarrow Pic^0 C^\sim$ is given by $\mathcal{F} \mapsto \pi^*(\mathcal{F})^{**}$.

Theorem 1.4: $\mathcal{F} \in \overline{Pic_r^0 C} \iff \exists \mathcal{L} \in Pic^0 C$ such that $\mathcal{F} \otimes \mathcal{L} \in \overline{K}_r$.

Proof: Let $\mathcal{G}$ be an $\mathcal{O}_{C \times Spec k[[t]]}$ -module such that $(\mathcal{G}/t\mathcal{G}) \approx \mathcal{F}$ and $\mathcal{G}$ restricted to the generic point of $Spec k[[t]]$ is generated by $r + 1$ elements

⁴ M_τ of [PS] is analogous to this.

⁵ [PS, Remark 3] is analogous to this.

⁶ (1.2), (1.3) (1.5) answer questions from Verdieri in private talk (Pondicherry, Jan. '89) who pointed out that although these results follow from the rest of this section, they deserve to be mentioned independantly.

modulo $\mathcal{C}$. We have a diagram

$$\begin{array}{ccc}
h & : & \text{Spec } k[[t]] \longrightarrow \text{Pic}^0 C, \\
h_0 & : & \text{Spec } k(t) \longrightarrow \text{Pic}_r^0 C \\
& & \downarrow \pi_r^* \\
\text{Pic}^0 C & \xrightarrow{\pi^*} & \text{Pic}^0 C^\sim
\end{array}$$

with the generic point of $\text{im}(h)$ given by h_0 . Since $\text{Pic}^0 C^\sim$ is complete, $\pi_r^* oh_0$ lifts to a map $p_0 : \text{Spec } k[[t]] \longrightarrow \text{Pic}^0 C^\sim$, and since π^* is smooth and surjective, $\text{im}(p_0)$ lifts to a curve in $\text{Pic}^0 C$, defining a line bundle $\mathcal{K}^{-1}$ on $C \times \text{Spec } k[[t]]$. Let $\mathcal{P} = \mathcal{K} \otimes \mathcal{G}$. Then $(\mathcal{P}/t\mathcal{P}) \approx \mathcal{L} \otimes \mathcal{F}$ where $\mathcal{L} = (\mathcal{K}/t\mathcal{K})$ and $\mathcal{P}$ restricted to the generic point of $\text{Spec } k[[t]]$ is in K_r . This proves one direction, and since the $\text{Pic}^0 C$ -orbit of K_r is contained in $\overline{\text{Pic}_r^0 C}$ by definition, it proves the result. $\parallel$

Finally, since $\pi_*(\text{Pic}^{-\delta} C^\sim) \subseteq B(\overline{\text{Pic}_r^0 C}) \ \forall r \in \{0, \dots, v_0 - 2\}$, we have

Lemma 1.5: $(\text{Pic}^0 C)^\sim$ is connected. ⁷ $\parallel$

⁷This is [PS, Th. 2].

2: $\mathcal{M} = \mathcal{C}$:

If ⁸ $\mathcal{M} = \mathcal{C}$ then $\overline{Pic_r^0 C}$ is a component of $(Pic^0 C)^\circ$ for $r = 0, \dots, v_0 - 2$, and we look at the intersections of components. (From this section on readers are referred to [J], [PS] for proofs.)

Theorem 2.1: If $\mathcal{M} = \mathcal{C}$ then $\overline{K_r} = E(\mathcal{C}, \delta - r - 1)$. ||

Consequently if $\mathcal{M} = \mathcal{C}$ then $\overline{Pic_r^0 C} = \text{Orb}_{Pic^0 C} E(\mathcal{C}, \delta - r - 1)$.

Theorem 2.3: If $\mathcal{M} = \mathcal{C}$ then $\overline{Pic_r^0 C}$ is an irreducible component of $(Pic^0 C)^\circ$ for $r = 0, \dots, v_0 - 2$. ||

Lemma 2.4: If $\mathcal{M} = \mathcal{C}$ then

(i) $B(\overline{Pic_r^0 C}) = \pi'_*(\overline{Pic_r^{-1} C'})$.

(ii) $\overline{Pic_r^0 C} \cap \overline{Pic_{r+1}^0 C} = \phi$.

(iii) $\overline{Pic_r^0 C} \cap \overline{Pic_{r+j}^0 C} = \pi_*^j(\overline{Pic_r^{-j} C_j})$ for $r = 1, \dots, v_0 - 2$ and $j = 1, \dots, v_0 - 2 - r$. ||

We summarize the section in a couple of graphs. Arrows indicate that the object at source of arrow is contained in the object at the tip of the arrow.

⁸The curve type $\mathcal{M} = \mathcal{C}$ of our notation is the type $\Gamma_{m,1,1}$ of [PS].

$$\begin{array}{ccccccc}
Pic_0^0 C & & & & & & \\
& \nwarrow & & & & & \\
& & Pic_0^{-1} C' & & & & \\
& & & & & & \\
Pic_1^0 C & & & & & & \\
& \nwarrow & & & & & \\
& & Pic_1^{-1} C' & & \dots & & \\
\dots & & \dots & \dots & \dots & \dots & \\
\dots & & \dots & \dots & \dots & \dots & \\
\dots & & \dots & \dots & \dots & \dots & \\
Pic_r^0 C & & \dots & \dots & \dots & & \\
& \nwarrow & & \dots & \dots & & \\
& & Pic_r^{-1} C' & & \dots & & \\
& & & & \dots & & \\
Pic_{r+1}^0 C & & \dots & \dots & \dots & & \\
& \nwarrow & & & \dots & & \\
\dots & & \dots & \dots & \dots & \dots & \\
\dots & & \dots & \dots & \dots & \dots & \\
\dots & & \dots & \dots & \dots & \dots & \\
Pic_{v_0-3}^0 C & & & & & & \\
& \nwarrow & & & & & \\
& & Pic_{v_0-3}^{-1} C & & & & \\
& & & & & & \\
Pic_{v_0-2}^0 C & & & & & &
\end{array}$$

$$\begin{array}{ccccccc}
\overline{Pic_0^0 C} & & & & & & \\
& \swarrow & & & & & \\
& & \overline{Pic_0^{-1} C'} & & & & \\
& \swarrow & & \dots & & & \\
\overline{Pic_1^0 C} & & & \dots & & & \\
& \swarrow & & & & & \\
\dots & \dots & \dots & \dots & \dots & & \\
\dots & \dots & \dots & \dots & \dots & \dots & \\
& \swarrow & & \dots & \dots & \swarrow & \\
\overline{Pic_r^0 C} & & & \dots & \dots & & \overline{Pic^{2-\delta} C^{\delta-2}} \\
& \swarrow & & \dots & \dots & \swarrow & \\
& & \overline{Pic_r^{-1} C'} & & \dots & & \overline{Pic^{1-\delta} C^{\delta-1}} \\
& \swarrow & & \dots & \dots & \swarrow & \\
\overline{Pic_{r+1}^0 C} & & & \dots & \dots & & \overline{Pic_1^{2-\delta} C^{\delta-2}} \\
& \swarrow & & \dots & \dots & \swarrow & \\
\dots & \dots & \dots & \dots & \dots & \dots & \\
\dots & \dots & \dots & \dots & \dots & & \\
& \swarrow & & & & & \\
\overline{Pic_{v_0-3}^0 C} & & & \overline{Pic_{v_0-3}^{-1} C'} & & & \\
& \swarrow & & & & & \\
& & & & & & \\
\overline{Pic_{v_0-2}^0 C} & & & & & &
\end{array}$$

3: $\mathcal{M}' = \mathcal{C}'$, C Gorenstein:

$\mathcal{O}$ is Gorenstein if and only if every torsion free $\mathcal{O}$ -module is reflexive, i.e, if and only if $\Gamma = \{j \mid v_0 - 1 - j \notin \Gamma\}$. So for all $\mathcal{F} \in \text{Pic}^0 C$, we have $\text{End}(\mathcal{F}) = \mathcal{O}_C \iff \mathcal{F}_Q \approx \mathcal{O}, \iff \mathcal{F} \in \overline{\text{Pic}^0 C}$. In particular therefore the components of $(\text{Pic}^0 C)^\perp$ other than $\overline{\text{Pic}^0 C}$, if any, are components of $\pi'_*(\text{Pic}^{-1} C')^\perp$. If C is locally planar, we know that $\overline{\text{Pic}^0 C} = (\text{Pic}^0 C)^\perp$ and $\pi'_*(\text{Pic}^{-1} C')^\perp$ is all of $B(\overline{\text{Pic}^0 C})$, contributing no components to $(\text{Pic}^0 C)^\perp$. On the other hand, if $\mathcal{M}' = \mathcal{C}'$ and if C is not locally planar⁹ then $\Gamma = (k_1, \dots, 2k_1 - 2)\mathbf{N}$, $k_1 \geq 3$.

Theorem 3.1: Let C be Gorenstein, and not locally planar. If $\mathcal{M}' = \mathcal{C}'$, then

- (i) $(\text{Pic}^0 C)^\perp = \text{Pic}^0 C \cup \pi'_*(\text{Pic}^{-1} C')^\perp$, and components of $(\text{Pic}^0 C)^\perp$, other than $\overline{\text{Pic}^0 C}$, are precisely $\{\pi'_*(\text{Pic}_r^{-1} C') \mid r = 1, \dots, \delta - 2\}$.
- (ii) If $\mathcal{F} \in \overline{\text{Pic}^0 C}$, and if $F \approx \mathcal{F}_Q$ is such that $F \in E(\mathcal{C}, \delta)$, then $\forall r \geq 1$, we have $\text{ord}_t F = r \implies \mathcal{F} \in \pi'_*(\overline{\text{Pic}_{r-1}^{-1} C'})$. Conversely, for all $\mathcal{F} \in \pi'_*(\overline{\text{Pic}_{r-1}^{-1} C'})$, $\mathcal{F} \in \overline{\text{Pic}^0 C} \implies \exists F \approx \mathcal{O}_Q$ such that $\mathcal{C} \subseteq F \subseteq \mathcal{O}^\sim$, with $\text{ord}_t F = r$, $F_1 \subseteq t^{k_1} \mathcal{O}^\sim$ and $\text{rk}(F_1/t^{k_1+r} \mathcal{O}^\sim) = r - 1$. ||

Example 1: Let $\mathcal{O}$ be generated by monomials in some local parameter t , and let $F = (t^2, t^{k_1})\mathcal{O} + \mathcal{C}$. The only possible deformation of F is given by either $(t^2 + \beta)\mathcal{O}$ when k_1 is even or $(t^2 + \beta t + \beta^2)\mathcal{O}$ when k_1 is divisible by 3, and there is no deformation if $k_1 \equiv \pm 1$ modulo 6. ◇

So the reverse implications of (ii) are not true. On the other hand

⁹ This is $\Gamma_{m, m-1}$ of [PS].

Example 2: For $\mathcal{O} = k[[t^4, t^5, t^6]]$ we have $Filt(\mathcal{C}, \delta) = \overline{K}$: if $ord_t F = 1$ then $F = t\mathcal{O}'$; if $ord_t F = 2$ and $F \neq t^2\mathcal{O}''$ then $F = (t^2, ut^4) + \mathcal{C}$ for $u \in \mathcal{O}^{\sim*}$, and $\partial_\beta = t^2 + \beta u$ will do; if $ord_t F = 3$ then F is in the closure of the Picard orbit of $(t^3, t^4, t^5) + \mathcal{C} = t^{-1}\mathcal{M}$, and $ord_t F = 4$ if and only if $F = \mathcal{O}^\sim$. In any case F has the desired deformation. $\diamond$

4: $Rk(\mathcal{M}/\mathcal{C}) = 1$, $\mathcal{M}' = \mathcal{C}'$:

From (0.13), if $rk(\mathcal{M}/\mathcal{C}) = 1$, then $Filt(\mathcal{C}, \delta) = \overline{K}$. By definition each $\overline{Pic_r^0 \mathcal{C}}$ contains at least one component of $(Pic^0 \mathcal{C})^\sim$ for $j \in \{1, \dots, k_1 - 1\}$. Since $(Pic^0 \mathcal{C})^\sim = \bigcup \{ \overline{Pic_r^0 \mathcal{C}} \mid r = 1, \dots, k_1 - 1 \}$, we need to find components of $\overline{Pic_r^0 \mathcal{C}}$ and to see when these are actually components of $(Pic^0 \mathcal{C})^\sim$. If $Rk(\mathcal{M}/\mathcal{C}) = 1$ and $\mathcal{M}' = \mathcal{C}'$ ¹⁰ then we have $\overline{Pic_0^0 \mathcal{C}} = \phi$, $Pic_1^0 \mathcal{C} = Pic^0 \mathcal{C}$, $Pic_2^0 \mathcal{C} = \{ \mathcal{F} \mid \mathcal{F}_Q = \mathcal{O} + f_1 \mathcal{O}, ord_t f_1 \geq 2 \}$, and $Pic_{k_1}^0 \mathcal{C} = orb_{Pic^0 \mathcal{C}} \mathcal{W}_\mathcal{C}^0$. Let $\mathcal{F} \in Pic_r^0 \mathcal{C}$ and $F \approx \mathcal{F}_Q$ such that $u\mathcal{O} \subseteq F \subseteq \mathcal{O}^\sim$ and $u \in \mathcal{O}^{\sim*}$, so that $rk(F/\mathcal{C}) = r + 1$. Let $\{f_0, f_1, \dots, f_r\}$ be a basis of F , $f_i = \sum \{a_{ij} t^j \mid j = 0, \dots, k_1 + 1, a_{ij} \in k\}$, and, say, $a_{00} \neq 0$. Then we must have $rk[(f_0, \dots, f_r)\mathcal{O} + \mathcal{C}]/\mathcal{C} = r + 1$,

$$\begin{aligned} \implies & \text{rank of } M_F, \text{ the coefficient matrix of } (F/\mathcal{C}), \text{ is } r + 1. \\ \implies & \text{every } (r + 2) \times (r + 2) \text{ minor of } M_F \text{ must vanish.} \end{aligned}$$

Therefore $\overline{Pic_r^0 \mathcal{C}} = \overline{P_{r,1}} \cup \overline{P_{r,2}}$, where

$$P_{r,1} = \{ \mathcal{F} \in Pic_r^0 \mathcal{C} \mid \exists F \approx \mathcal{F}_Q \text{ such that } F \in E(\mathcal{C}, \delta - r - 1), \mathcal{O} \subseteq F, \text{ and } ord_t F_1 = 1 \}$$

and

$$P_{r,2} = \{ \mathcal{F} \in Pic_r^0 \mathcal{C} \mid \exists F \approx \mathcal{F}_Q \text{ such that } F \in E(\mathcal{C}, \delta - r - 1), \mathcal{O} \subseteq F, \text{ and } ord_t F_1 \geq 2 \}.$$

It remains to see whether or not $\overline{P_{r,i}}$ are components of $Pic^0 \mathcal{C}$ for $r = 1, \dots, k_1 - 1$. Since $P_{r,1} = \pi'_* Pic_{r-2}^{-1} \mathcal{C}'$ for $r = 2, \dots, k_1$, $\overline{P_{r,1}}$ is not contained in $P_{r',1}$ for $r \neq r'$. Hence proving that $P_{r,i}$ are, indeed, components of $(Pic^0 \mathcal{C})^\sim$, is now reduced to the question of looking at $B(\overline{P_{r,2}})$.

Theorem 4.1: $\mathcal{F} \in \overline{P_{r,2}}$ if and only if the following conditions hold:

- (i) $\exists F$ such that $\mathcal{F}_Q \approx F$, $\mathcal{C} \subseteq F \subseteq \mathcal{O}^\sim$, $rk(F/\mathcal{C}) = r + 1$, and
- (ii) either $ord_t F = 1$ and $F \supseteq t^{k_1+1} \mathcal{O}^\sim$, or $ord_t F \geq 2$, where $F \supseteq F_1 \supseteq \dots \supseteq F_r \supseteq \mathcal{C}$ is a filtration with the following properties:

¹⁰ If $\mathcal{C}$ is a monomial singularity curve then this is $\Gamma_{m,1,2}$ of [PS].

$rk(F_i/F_{i+1}) = 1$, $ord_t F_{i+1} > ord_t F_i$, and $F_r \subseteq t^{k_1} \mathcal{O}^\sim$. ||

Corollary 4.1.1: $P_{i,1} \subseteq \overline{P_{i-1,2}}$ for all $i < k_1$. ||

Corollary 4.1.2: $\mathcal{F} \in \overline{P_{r,2}} \cap \overline{P_{r+1,2}} \iff \exists F \approx \mathcal{F}_Q$ such that $\mathcal{C} \subseteq F \subseteq t\mathcal{O}^\sim$, $rk(F/\mathcal{C}) = r+1$ and $F_1 \subseteq t^3 \mathcal{O}^\sim$. ||

For $i = 1, \dots, k_1 - 1$, $\exists \mathcal{F} \in P_{i,2}$ such that $\mathcal{F}_Q \approx \sum \{t^j \mathcal{F} \mid j = 0, 2, \dots, i\}$. Since $\mathcal{F} \notin P_{i',2} \forall i \neq i'$, this proves

Theorem 4.2: $(Pic^0 C)^\sim = \bigcup \{P_{i,2} \mid i = 1, \dots, k_1 - 1\}$ and $\overline{P_{i,2}}$ is a component of $(Pic^0 C)^\sim$ for $i = 1, \dots, k_1 - 1$. ||

References

- [AIK] A. Altman, A. Iarrobino and S. Kleiman, Irreducibility of the compactified Jacobian, in "Proceedings, Nordic Summer School NAVF, Oslo, Aug. 5-25, 1976," Noordhoff, Groningen, 1977.
- [AK1] A. Altman and S. Kleiman, Compactifying the Picard Scheme, Advances In Mathematics 35, 50-112 (1980).
- [AK] A. Altman and S. Kleiman, Compactifying the Picard Scheme, II, Amer. J. Math. 101 (1979) ("Zariski volume"), 10-41.
- [D] C. D'Souza, "Compactification of Generalized Jacobians," Thesis, Tata Institute, Bombay, 1973.
- [HK] Herzog and Kunz, Die Wertehalbgruppe eines Lokalen Ringes der Dimension I, Springer Verlag, 1971.
- [J] _____, Thesis, Brandeis University, 1986.
- [J'] _____, Boundary of Picard Group of a Singular Curve, preprint.
- [K] H. Kleppe, Picard scheme of a curve and its compactification. Thesis, M.I.T., 1981.
- [MM] A. Mayor and D. Mumford, Further Comments on Boundary Points, Amer. Math. Soc. Summer Institute, Woods Hole, Mass., 1964.
- [PS] G. Pfister and J.H.M. Steenbrink, Reduced Hilbert Scheme for Irreducible Curve Singularities, to appear, Journal of Pure and Applied Algebra.

- [R] C. Rego, The Compactified Jacobian, Ann. Scient. Ec. Norm. Sup., 4⁴ serie, t. 13, 1980, p. 211 a 224.
- [SK] S. Kleiman: The structure of the compactified Jacobian : a review and an announcement; seminari di geometria, 1982-83; Universita delgi Studi di Bologna, Dipartimento Mathematica, 1984.